

Fusion Alpha Performance in Advanced Scenario Plasmas based on Reversed Central Magnetic Shear

T. Kurki-Suonio¹, O. Asunta¹, V. Tulkki¹, S. Sipilä¹, R. Salomaa¹, T. Tala²,
and JET-EFDA contributors*

¹ *Euratom-Tekes Association, Helsinki University of Technology*

² *Euratom-Tekes Association, VTT Energy, Finland*

Introduction. The successful operation of a fusion reactor relies on sufficient confinement of the fusion-born alpha particles. The 3.5 MeV fusion alphas can have orbit widths of the order of the size of the plasma column, and their confinement generally requires a very large plasma current. Therefore the alpha particle confinement becomes particularly worrisome for plasmas with a so-called internal transport barrier (ITB) [1], which is facilitated by very low or, preferably, reversed central magnetic shear [2]. For brevity we shall refer to such a profile simply as *reversed*, and it implies very low plasma current over a significant part of the plasma. The orbit widths are generally inversely proportional to the poloidal magnetic field, the strength of which scales with the plasma current. Therefore, in the presence of a current hole or very low central plasma current, the confinement of fusion-born alphas becomes a major concern.

We study the confinement and heating profile of fusion alphas both in a normal JET H-mode plasma (shot 52009) with a monotonic q -profile and in an ITB plasma (shot 51976) with a reversed q -profile and a current hole (CH), see Fig. 1(a). The work is carried out using the Monte Carlo -based guiding-center-following code ASCOT [3] which naturally and accurately resolves all possible orbit topologies as well as the transitions between them [4]. This work complements the recent Fokker-Planck analysis [5] not only by using a completely different method but also by allowing the experimental flux surface structure shown in Fig. 1(b) which, in the center of an ITB plasma, does not easily render itself to simple parametrization.

Orbits of 3.5 MeV alphas with normal and reversed q -profile. Figure 2 shows drift orbits for a 3.5 MeV alpha launched at various locations along the midplane for shots 51976 and 52009. The orbit topologies are dominated by potatoes and passing orbits, with bananas existing only in a very limited region. The orbit width reaches a local minimum when the starting point is moved outwards from the plasma center and the poloidal field reaches a critical value at which the particle's own poloidal velocity at the midplane is cancelled by the gradient drift. For fusion alphas in JET geometry this field is given by [6] $B_p = mv/R_0e \approx 90$ mT. As such a location is reached, the spatial extent of the particles' orbits in the poloidal plane is reduced to that of their gyromotion, not resolved by the present technique. These orbits are called *stagnation*




Figure 3: *The plasma profiles for an ITB (a) and H-mode (b) discharges. Also shown is the local fusion alpha production rate, particles per second.*




Figure 4: *The power (W) deposited to plasma by fusion alphas. The values are normalized to maximum value.*

test alphas. During the simulation, the change in particle energy is recorded, along with its location in the poloidal cross-section, yielding the power deposition profile. The radial profile of the alpha heating power, in watts, is illustrated in Fig. 4. To facilitate the comparison, the profiles are normalized to their maximum value. As function of ρ the H-mode shot yields more distributed heating, mostly due to the radically different plasma profiles. This was verified by repeating the H-mode simulation using the plasma profiles of the ITB shot. Also, the physical volume corresponding to ρ is quite different in these two shots.

The power density profile in the poloidal cross-section is displayed in Fig. 5, again normalized to its maximum value. Any poloidal differences have to be attributed to the magnetic structures because plasma profiles are flux functions. In the H-mode, shot 52009, the heating strongly peaks in the center, favoring only slightly the LFS due to the trapped orbits. For the shot 51976 with an ITB, the heating shows structure around the two competing magnetic axes inside the current hole. The differences in the LFS/HFS asymmetry between the two shots reflect the orbit topologies: H-mode favors the LFS due to the trapped orbits that exist only there, while with a CH the presence of potato orbits even on the HFS keeps the heating power more uniform. Neither H-mode nor ITB shot exhibits any poloidally localized heating maximum due to stagnation orbits, as was observed in an earlier study [6]. Also, from this figure it is clear that the physical area over which the alphas deposit energy is larger for the ITB shot.

Conclusions. In agreement with a recent study [11], advanced scenarios based on current hole were found to provide a broad, uniform alpha heating profile. This allowed the same total heating with 20% smaller plasma current. Next the simulations will be refined with a particle loading better adapted for current holes.




Figure 5: Poloidal distribution of the alpha heating power density (W/m^3) for a) an ITB discharge, and b) an H-mode discharge. The power is normalized to its maximum value.

* See annex of J. Pamela *et al.* Proc. 20th Int. Conf. Fusion Energy (Vilamoura, 2004) OV/1-2 (Vienna:IAEA)

Acknowledgements: The computing facilities of CSC were used in this work. This work, supported by the European Communities, under the contract of Association between Association Euratom/Tekes, was carried out within the framework of the European Fusion Development Agreement. The views and opinions expressed herein do not necessarily reflect those of the European Commission.

1. X. Litaudon *et al.*, Plasma Phys. Control. Fusion **41** No 3A (March 1999) A733.
2. X. Litaudon *et al.*, Plasma Phys. Control. Fusion **48** (2006) A1.
3. J.A. Heikkinen *et al.*, Journal Comput. Phys. **173** (2001) 527.
4. L.-G.Eriksson and F. Porcelli, Plasma Phys. Control. Fusion **43** (2001) R145.
5. V. Yavorskij *et al.*, Nuclear Fusion **43** (2003) 1077.
6. T. Kurki-Suonio *et al.*, Fusion Alpha Performance in Advanced Scenario Plasmas based on Reversed Central Magnetic Shear, submitted to Plasma Phys. Contr. Fusion.
7. L.M. Hively, G.H. Miley, J.A. Rome, Nuclear Fusion **21**, No. 11 (1981) 1431.
8. N.C. Hawkes *et al.*, Plasma Phys. Control. Fusion **47** (2005) 1475.
9. H.H. Duong *et al.*, Physical Review Letters **75**, No. 5 (1995) 846.
10. H.-S. Bosch and G.M. Hale, Nuclear Fusion **32** (1992) 611.
11. M. Schneider *et al.*, Plasma Phys. Control. Fusion **47** (2005) 2087.