

Application of laser-accelerated high-energy protons for isochoric heating of matter

P. Antici^{1,2}, J. Fuchs¹, S. Atzeni², A. Benuzzi¹, E. Brambrink¹, M. Esposito¹, M. Koenig¹,
A. Ravasio¹, J. Schreiber³, A. Schiavi², P. Audebert¹

¹*Laboratoire pour l'Utilisation des Lasers Intenses, UMR 7605 CNRS-CEA-École Polytechnique-Univ. Paris VI, 91128 Palaiseau, France*

²*Dipartimento di Energetica ed INFM, Università di Roma « La Sapienza », Via Scarpa 14-16, 00165 Roma, Italy*

³*Max-Planck-Institut für Quantenoptik and Physik Department, Ludwig-Maximilians-Universität München, 85748, Garching, Germany*

The possibility of producing, in a controlled way, matter at solid density (1-10 g/cm³) while maintaining it at a high temperature (1-100 eV) has been for a long time a desired goal. Studying matter in these conditions would provide an array of information useful for a variety of problems [1] that range from fundamental physics (dense and relatively cold matter at the frontier of plasma physics) to astrophysics (matter in the planetary cores) through applications like ICF (stopping power of particles in warm dense matter).

To achieve this, energy has to be deposited in a short time scale before heating-induced expansion decreases the density of the heated matter, i.e. isochorically. The heated material is then inertially confined. Up to now, attempts to achieve isochoric heating have been pursued using either short-pulses lasers or short-duration laser-produced x-rays (through inner transitions induced in solids by laser-accelerated hot electrons). Optical frequencies are very limited in their penetration of solid matter, typically some nm of skin depth in Al for 1 μ m light. Therefore only a surface layer is directly heated [2] while the inner part of the sample gets heated on a much longer time scale through heat conduction accompanied by thermal expansion. This does not allow maintaining a high density and temperature. Therefore, high intensity laser pulses incident on a thin foil cannot create an uniform, hot dense plasma.

In comparison, ions have a much higher penetration depth, allowing *volumetric heating*; for example the Bragg peak in Al is at a depth of 5 μ m for a 0.5 MeV proton and 200 μ m for a 5 MeV proton. Ions deposit energy directly and deeply in matter and can create isochoric heating provided that the energy deposition time is shorter than the heating-induced expansion time. In typical large ion accelerators such as GSI, much higher ion energies can be produced




Figure 2: Separation of the OTR signal induced by fast electrons from the proton heating signal a) in time using a fast streak camera b) in space, using a tilted target.

During the experiment, we have explored a range of parameters by varying the thickness of the sample heated foil, using targets of different Z like aluminum or gold, varying the heated temperature by moving the sample out of focus and using curved targets.

Figure 3 shows the time-resolved 1D optical emission from the rear-surface of various proton-heated targets. One can see that the heating decreases as the target thickness increases and that heating increases with the Z of the target, as expected. This, since the stopping power of the protons increases with higher Z targets.

Experimental data are compared to simulations using a 2D hydrocode (DUED) [5] and having as input the known accelerated proton energy distribution. This distribution is made out of two temperatures, one for the low energy protons ($T=0.9\text{MeV}$) and one for the high energy protons ($T=5.4\text{MeV}$). For both, the number of protons decreases exponentially, the high energy proton distribution having a higher temperature, but a smaller number of protons (i.e. there are 2 orders of magnitude between the number of protons at 0.5 and 5 MeV). The main contribution of the heating is predominantly due to the low energy protons. Figure 4 shows how the target can be heated in a few picoseconds to temperatures of a few eV. Moreover one can clearly see the initial isochoric heating and, after some tens of picoseconds, the adiabatic expansion. Also a radial profile of heating is detectable. On the back side of the target, where no protons are arriving, the adiabatic expansion is visible.

Concerning curved targets [4], we were able to better concentrate protons on the target, but did not see a substantial increase of the heating. It seems hence better to focus the laser with its full intensity on a flat target than to irradiate a larger curved spot which is necessary to focus the protons but causes a too large decrease in the number of accelerated protons.




Figure 3: Time-resolved 1D images of the rear-surface of a proton-heated foil. Time goes up.




Figure 4: DUED Simulation for a 25 μm aluminum foil heated by protons, 20ps after the irradiation starts. a) shows the 2D temperature map whereas in b) is shown in abscissa the longitudinal target depth, in ordinate the mean temperature around the central point ($\pm 15\mu\text{m}$). The Bragg peak one can see at 20 ps is mainly due to protons of about 0.5MeV, which cross the distance between the two targets of about 200 μm in 20ps.

The results show that accelerated protons can indeed be used to achieve heating of solid targets. This method is promising as a way either to measure the equations of state of materials in poorly known conditions (a few eV, solid density) or as a way to measure the stopping power of higher Z ions (other than protons) which is not well known.

This work was supported by DAAD, grant E1127 from Région Ile-de-France and University of Nevada, Reno for target support (G.Korgan, S.Malekos, N.Le Galloudec, T.Cowan).

References:

- [1] <http://www-hasylab.desy.de/conferences/WDM/content-3.html#wdm>
- [2] P. Audebert *et al.*, Phys. Rev. Lett. **89**, 265001 (2002).
- [3] R. Snavely *et al.*, Phys. Rev. Lett. **85**, 2945 (2000); S. C. Wilks *et al.*, Phys. Plasmas **8**, 542 (2001) ; S. Hatchett *et al.*, Phys. Plasmas **7**, 2076 (2000) ; M. Hegelich *et al.*, Phys. Rev. Lett. **89**, 085002 (2002).
- [4] P. K.Patel *et al.*, Phys. Rev. Lett. **91**, 125004 (2003)
- [5] S. Atzeni *et al.*, Comput.Phys.Commun doi:10.1016/J.cpc.2005.03.03